



## Original Article


## Phase shift, amplification, oscillation and attenuation of solitons in nonlinear optics

Weitian Yu<sup>a</sup>, Qin Zhou<sup>b,\*</sup>, Mohammad Mirzazadeh<sup>c</sup>, Wenjun Liu<sup>a,d,\*</sup>, Anjan Biswas<sup>e,f</sup><sup>a</sup> State Key Laboratory of Information Photonics and Optical Communications, School of Science, P.O. Box 122, Beijing University of Posts and Telecommunications, Beijing 100876, China<sup>b</sup> School of Electronics and Information Engineering, Wuhan Donghu University, Wuhan 430212, China<sup>c</sup> Department of Engineering Sciences, Faculty of Technology and Engineering, East of Guilan, University of Guilan, P.C. 44891-63157, Rudsar-Vajargah, Iran<sup>d</sup> Beijing National Laboratory for Condensed Matter Physics, Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China<sup>e</sup> Department of Physics, Chemistry and Mathematics, Alabama A&M University, Normal, AL 35762, USA<sup>f</sup> Department of Mathematics and Statistics, Tshwane University of Technology, Pretoria 0008, South Africa

## HIGHLIGHTS

- Effects of the reciprocal of group velocity on solitons were discussed.
- Energy exchange of solitons occurred during the phase shift.
- Solitons in a bound state were amplified or attenuated.
- Parabolic soliton interactions were analysed to decrease the interactions.
- Parabolic solitons can be reduced to dromion-like structures.

## GRAPHICAL ABSTRACT



Interactions between solitons can be controlled through adjusting the phase shift of solitons.

## ARTICLE INFO

## Article history:

Received 14 July 2018

Revised 5 September 2018

Accepted 6 September 2018

Available online 7 September 2018

## Keywords:

Solitons  
Amplification  
Oscillation  
Attenuation  
vcNLS

## ABSTRACT

In nonlinear optics, the soliton transmission in different forms can be described with the use of nonlinear Schrödinger (NLS) equations. Here, the soliton transmission is investigated by solving the NLS equation with the reciprocal of the group velocity  $\beta_1(z)$ , the group velocity dispersion coefficient  $\beta_2(z)$  and nonlinear coefficient  $\gamma(z)$ . Two-soliton solutions for the NLS equation are obtained through the Hirota method. According to the solutions obtained,  $\beta_1(z)$  and  $\gamma(z)$  with different function forms are taken to study the characteristics of solitons. The effect of the phase shift on the soliton interaction is discussed, and the non-oscillating soliton amplification, which is transmitted in a bound state, is explored. Parabolic solitons with oscillations are analysed. Moreover, parabolic solitons can be reduced to dromion-like structures. Results indicate that the transmission of solitons can be adjusted with the group velocity dispersion and Kerr nonlinearity coefficients. The phase shift, amplification, oscillation and attenuation of solitons can also be controlled by other related parameters. This work accomplishes the theoretical study of transmission characteristics of optical solitons in spatially dependent inhomogeneous optical fibres. The conclusions of this research have theoretical guidance for the research of optical amplifier, all-optical switches and mode-locked lasers.

© 2018 Production and hosting by Elsevier B.V. on behalf of Cairo University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer review under responsibility of Cairo University.

\* Corresponding authors.

E-mail addresses: [qinzhou@whu.edu.cn](mailto:qinzhou@whu.edu.cn) (Q. Zhou), [jungliu@bupt.edu.cn](mailto:jungliu@bupt.edu.cn) (W. Liu).<https://doi.org/10.1016/j.jare.2018.09.001>

2090-1232/© 2018 Production and hosting by Elsevier B.V. on behalf of Cairo University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Solitons have been investigated in such fields as mathematics and physics [1–6]. They can propagate in a long distance without changes in their waveform, velocity and amplitude [7–10]. The soliton phenomena are closely related to nonlinear evolution equation models [11–17]. Solitons are also applied in particle physics, fluid mechanics, Bose-Einstein condensation and nonlinear optics [18–20].

Some researchers have studied solitons by solving the nonlinear evolution equations and analysing their soliton solutions [21–25]. As one of the classic nonlinear evolution equations, the nonlinear Schrödinger (NLS) equation can be solved to obtain soliton solution, and has been widely investigated by using different methods [26–32]. With the differential quadrature method, the dynamic problems constructed by the NLS equation have been analysed [26]. The existence and stability of the standing wave solutions for the NLS equation in  $n$ -dimensional space have been studied [27]. Using the generalized exponential rational function, a new method to solve the exact special solutions of the NLS equation has been proposed [28]. The unified method has been used to acquire optical soliton solutions of the NLS equation [29]. Moreover, the stability of full dimensional KAM tori for the NLS equation has been proved [30]. Local Cauchy theory for the NLS equation has been discussed [31], and nonlinear instability of half-solitons has been analysed [32].

However, the soliton transmission process can be simulated more accurately with the variable coefficient NLS (vcNLS) equation, when the transmission medium or boundary condition is not uniform [33–40]. For the vcNLS equation, dynamics of solitons have been explored [33], and soliton interactions have been discussed [34,35]. In addition, the breather-to-soliton transitions for the vcNLS equation have been found [36], and nonautonomous multi-peak solitons have been obtained [37].

The different transmission characteristics of solitons can be obtained by solving this vcNLS equation,

$$i \frac{\partial u}{\partial z} + i\beta_1(z) \frac{\partial u}{\partial t} + \beta_2(z) \frac{\partial^2 u}{\partial t^2} + \gamma(z)|u|^2 u = 0, \quad (1)$$

where  $u(z, t)$  describes the temporal envelope of solitons;  $z$  and  $t$  represent the longitudinal coordinate and the time in the moving coordinate system, respectively; and  $\beta_1(z)$ ,  $\beta_2(z)$  and  $\gamma(z)$  are related to the reciprocal of the group velocity, group velocity dispersion (GVD) coefficient and the nonlinearity coefficient, respectively. If  $\beta_1(z) = 0$ , Eq. (1) will become the standard vcNLS equation.

Eq. (1) can be used to describe the transmission of solitons. The discussion of  $\beta_1(z)$  and  $\beta_2(z)$  is helpful to the development of dispersion management communication systems [38]. In addition, the study of  $\beta_1(z)$ ,  $\beta_2(z)$  and  $\gamma(z)$  is of significance to the experimental and engineering application of mode-locked fiber lasers and nonlinear optics. The rogue wave solutions for Eq. (1) have been solved with similarity transformation [39], and the properties of oscillating solitons for Eq. (1) have been analysed [40].

However, parallel solitons, parabolic solitons and dromion-like solitons obtained by taking different functions for  $\beta_1(z)$  and  $\gamma(z)$  have not been reported. The effect of the phase shift on the transmission of parallel solitons will be discussed. Here, the non-oscillating parallel soliton amplification, which transmits in bound states, will be studied. Parabolic solitons with oscillations will be analysed. The method of the parabolic soliton reducing to the dromion-like structure will be proposed by adjusting the corresponding parameters.

In Section 'Material and methods', the two-soliton solutions will be solved through use of the Hirota method. In Section 'Results and discussion', different soliton transmission characteristics are

analysed with the variable coefficients taking different functions. And in Section 'Conclusions', the conclusions will be derived.

## Material and methods

Firstly, the following independent variable transformation is introduced to solve the bilinear form according to the Hirota method [40],

$$u(z, t) = \frac{g(z, t)}{f(z, t)}, \quad (2)$$

where  $g(z, t)$  and  $f(z, t)$  are the complex function and real function, respectively. Then the bilinear form is obtained,

$$[iD_z + i\beta_1(z)D_t + \beta_2(z)D_t^2]g \cdot f = 0,$$

$$\beta_2(z)D_t^2 f \cdot f - \gamma(z)gg^* = 0.$$

where  $*$  is the complex conjugate. Moreover, the bilinear operators  $D_z$  and  $D_t$  are defined as follows

$$D_z^m D_t^n (G \cdot F) = \left( \frac{\partial}{\partial z} - \frac{\partial}{\partial z'} \right)^m \left( \frac{\partial}{\partial t} - \frac{\partial}{\partial t'} \right)^n G(z, t) F(z', t') \Big|_{z'=z, t'=t}.$$

According to the Hirota method, the bilinear form can be solved by the following power series expansions of  $g(z, t)$  and  $f(z, t)$ ,

$$g(z, t) = \varepsilon g_1(z, t) + \varepsilon^3 g_3(z, t) + \varepsilon^5 g_5(z, t) + \dots,$$

$$f(z, t) = 1 + \varepsilon^2 f_2(z, t) + \varepsilon^4 f_4(z, t) + \varepsilon^6 f_6(z, t) + \dots,$$

Here  $\varepsilon$  is a formal expansion parameter.

In order to obtain the two-soliton solutions, it can be assumed that

$$g(z, t) = \varepsilon g_1(z, t) + \varepsilon^3 g_3(z, t),$$

$$f(z, t) = 1 + \varepsilon^2 f_2(z, t) + \varepsilon^4 f_4(z, t),$$

where

$$g_1(z, t) = e^{Q_1(z, t)} + e^{Q_2(z, t)}, f_4(z, t) = n_5(z) e^{Q_1(z, t) + Q_2(z, t) + Q_1^*(z, t) + Q_2^*(z, t)},$$

$$g_3(z, t) = m_1(z) e^{Q_1(z, t) + Q_2(z, t) + Q_1^*(z, t)} + m_2(z) e^{Q_1(z, t) + Q_2(z, t) + Q_2^*(z, t)},$$

$$f_2(z, t) = n_1(z) e^{Q_1(z, t) + Q_1^*(z, t)} + n_2(z) e^{Q_2(z, t) + Q_2^*(z, t)} + n_3(z) e^{Q_1(z, t) + Q_2^*(z, t)} + n_4(z) e^{Q_2(z, t) + Q_1^*(z, t)}.$$

In addition,

$$Q_1(z, t) = k_{11}(z) + ik_{12}(z) + (w_{11} + iw_{12})t + \theta_{11} + i\theta_{12},$$

$$Q_2(z, t) = k_{21}(z) + ik_{22}(z) + (w_{21} + iw_{22})t + \theta_{21} + i\theta_{22}.$$

Finally, one can obtain the solutions,

$$m_1(z) = \frac{s_{12}\gamma(z)}{8w_{21}^2 s_{11}\beta_2(z)}, m_2(z) = \frac{s_{12}\gamma(z)}{8w_{21}^2 s_{11}\beta_1(z)},$$

$$\beta_2(z) = c\gamma(z), n_5(z) = \frac{[(w_{11} - w_{21})^2 + (w_{12} - w_{22})^2]^2}{64w_{11}^2 w_{21}^2 [(w_{11} + w_{21})^2 + (w_{12} - w_{22})^2]^2},$$

$$k_{11}(z) = - \int [w_{11}\beta_1(z) + 2w_{11}w_{12}\beta_2(z)] dz, n_1(z) = \frac{\gamma(z)}{8w_{11}^2 \beta_2(z)},$$

$$k_{21}(z) = - \int [w_{21}\beta_1(z) + 2w_{21}w_{22}\beta_2(z)] dz, n_2(z) = \frac{\gamma(z)}{2s_{21}\beta_2(z)},$$

$$k_{12}(z) = - \int [w_{12}\beta_1(z) - w_{21}^2\beta_2(z) + w_{12}^2\beta_2(z)]dz, n_3(z) = \frac{\gamma(z)}{2s_{11}\beta_2(z)},$$

$$k_{22}(z) = - \int [w_{22}\beta_1(z) - w_{21}^2\beta_2(z) + w_{22}^2\beta_2(z)]dz, n_4(z) = \frac{\gamma(z)}{8w_{21}^2\beta_2(z)},$$

and

$$s_{11} = (w_{11} + iw_{12} + w_{21} - iw_{22})^2, s_{12} = (w_{11} + iw_{12} - w_{21} - iw_{22})^2,$$

$$s_{21} = (w_{11} - iw_{12} + w_{21} + iw_{22})^2, s_{22} = (w_{11} - iw_{12} - w_{21} + iw_{22})^2.$$


### Results and discussion

By analysing the preceding two-soliton solutions, we find that if  $c = 1$ , then the GVD coefficient  $\beta_2(z)$  is equal to the nonlinearity coefficient  $\gamma(z)$ . Therefore, the different soliton transmission characteristics can be obtained by discussing the reciprocal of the group velocity  $\beta_1(z)$  and the nonlinearity coefficient  $\gamma(z)$ .

In order to obtain the effect of the phase shift on parallel soliton transmission, one takes  $\beta_1(z)$  as the Gauss function, and  $\beta_2(z) = \gamma(z) = 2\beta_1(z)$  as shown in Fig. 1. In Fig. 1(a) and (b), the phase shift of two solitons to the left influences the interaction

between two solitons. In order to keep two solitons from interacting with each other after the phase shift, one can adjust the parameter  $w_{11}$ . Comparing Fig. 1(b) with Fig. 1(c) or Fig. 1(a) with Fig. 1 (d), one can find that the direction of the phase shift is related to the signs of  $w_{12}$  and  $w_{22}$ . When  $w_{12}$  and  $w_{22}$  are positive, two solitons shift to the left, while two solitons move in opposite directions when they are negative. The larger the value of  $|w_{12}|$  or  $|w_{22}|$ , the greater is the phase shift distance of the soliton. Therefore, one can adjust the value of  $w_{22}$  as shown in Fig. 2 in order to separate the two interacting solitons after the phase shift. Moreover, the amplitude of the soliton increases as  $|w_{11}|$  or  $|w_{21}|$  increases.

To observe whether soliton energy exchange occurs during the phase shift, we can measure the amplitude of the soliton before and after the phase shift as shown in Fig. 2. In Fig. 2(a) and (b), the amplitudes of two solitons before and after the phase shift have obvious changes. However, the amplitudes of two solitons before and after the phase shift are almost the same in Fig. 2(c) and (d). Therefore, when two solitons are simultaneously shifted to the left, the energy transfer between solitons occurs. When two solitons are in opposite phase shifts, they remain constant in energy. Moreover, it can be seen from Figs. 1(c) and 2(c) that two solitons retain shape and amplitude after the collision except for a certain phase shift, which indicates that the collision is elastic. These analyses can help eliminate the interaction between solitons and change the direc-



**Fig. 1.** The phase shift has an effect on the transmission of parallel solitons. The parameters are  $\theta_{11} = -1, \theta_{12} = 2, \theta_{21} = -1, \theta_{22} = 2, \beta_1(z) = 2e^{-0.5z^2}$  and  $\gamma(z) = 2\beta_1(z)$  with (a):  $w_{11} = -0.5, w_{12} = 0.031, w_{21} = 0.5$  and  $w_{22} = 0.69$ ; (b):  $w_{11} = -0.78, w_{12} = 0.031, w_{21} = -0.5$  and  $w_{22} = 0.69$ ; (c):  $w_{11} = -0.78, w_{12} = -1.1, w_{21} = -0.5$  and  $w_{22} = 0.69$ ; (d):  $w_{11} = 0.5, w_{12} = 0.031, w_{21} = -0.5$  and  $w_{22} = 0.53$ .




Fig. 2. Amplitude comparison between two solitons at  $z = -11$  (red dotted line) and  $z = 11$  (green solid line) in (a): Fig. 1(a); (b): Fig. 1(b); (c): Fig. 1(c); (d): Fig. 1(d).

tion of the soliton transmission. This feature is useful for reducing the bit error rate, improving the quality of optical communication and studying all-optical switches.

Then, the effect of the phase shift on parallel solitons with the oscillation is discussed. In order to obtain the parallel solitons with oscillations,  $\beta_1(z) = 2e^{-5z^2} + \sin z$  and  $\gamma(z) = 4e^{-0.5z^2}$  are taken as Fig. 3 shows. In Fig. 3(a) and (b), one can find that the amplitude and phase shift of solitons can be changed by adjusting the parameters. The amplitude of the left soliton increases as  $|w_{11}|$  or  $|w_{21}|$  is increased. However, the amplitude of the right soliton increases with the decreasing of  $|w_{11}|$  or the increasing of  $|w_{21}|$ . The direction of the phase shift for two solitons is related to the sign of  $w_{12}$  and  $w_{22}$ . The phase shift distance increases as the values of  $|w_{12}|$  and  $|w_{22}|$  are increased. And when two solitons are in opposite phase shifts, elastic collision occurs during the phase shift. Because they have no change in amplitude and shape before and after the phase shift as shown in Fig. 4(a) and (b). From Fig. 3(c) and (d), one can see that not only are two solitons simultaneously shifted to the right, but also two solitons are shifted to the left at the same time. Further, in the process of the phase shift, the energy of two solitons is exchanged. Their amplitude changes before and after the phase shift, as shown in Fig. 4(c) and (d). These investigations can help weaken the energy exchange with the oscillatory parallel soliton during the phase shift. These studies can be helpful in the study of mode-locked fibre lasers.


The effect of the phase shift on the parallel soliton transmission and energy exchange was investigated. Next, we discuss the non-oscillating parallel soliton amplification, which transmits in bound states. In order to study the transmission, one takes  $\beta_1(z) = -\frac{0.55}{1+5z}$  and  $\gamma(z) = 4e^{-0.5z^2}$  as shown in Fig. 5. Near the  $z = 0$  point, it is mainly the sudden changes in  $\beta_1(z)$  and  $\gamma(z)$  that lead to change in the interaction between two solitons. As a result, two solitons have exchanged energy.

In Fig. 5(a), the left soliton amplitude increases and the right soliton amplitude decreases near the  $z = 0$  point. Energy is exchanged from the right soliton to the left. The situation in


Fig. 5(a) and (d) is just the opposite. In Fig. 5(b), the amplitudes of two solitons are reduced at the same time near the  $z = 0$  point, while the amplitudes of two solitons are amplified in Fig. 5(c). And in Fig. 5(a) and (d), the interaction between two solitons has also changed near the  $z = 0$  point. This is because the inverse function taken by  $\beta_1(z)$  is changed suddenly around the point of  $z = 0$ . However, before and after the point  $z = 0$  in Fig. 5(b) and (c), the interaction between two solitons does not change. Moreover, the amplitude is also affected by  $w_{ij}$  ( $i = 1, 2, j = 1, 2$ ).

Their changes in amplitude are shown in Fig. 6. The interaction between two solitons is inelastic. The energy is exchanged, and the amplitude changes between them without any outside influence. This can provide some help in the development of optical amplifiers.

To investigate the effect of  $\beta_1(z)$  and  $\gamma(z)$  on other types of soliton transmissions, one considers  $\beta_1(z) = -\cos(\arccos z)$  and  $\gamma(z) = \sin 2z$ . Then, a set of parabolic solitons with oscillations is obtained. In Fig. 7(a), two parabolic solitons maintain their original oscillation intensity due to the weak interaction. In Fig. 7(b), the interaction between two parabolic solitons causes the intensity of the inner parabolic soliton oscillations to become smooth. However, the interaction between parabolic solitons in Fig. 7(c) tends to smooth the intensity of the outer parabolic soliton oscillations. By analysing the parameters, one finds that  $w_{12}$  affects the soliton oscillation intensity of the inner side, and the inner soliton oscillates are exacerbated as  $|w_{12}|$  increases. The outer soliton oscillation intensity is controlled by  $w_{22}$  and exacerbated with the increasing of  $|w_{22}|$ . Moreover, the amplitude of the inner soliton is mainly controlled by  $|w_{11}|$ , while the amplitude of the outer soliton is mainly influenced by  $|w_{21}|$ . The amplitude of the inner and outer solitons is increased with increasing  $|w_{11}|$  and  $|w_{21}|$ , respectively. When  $|w_{11}| \rightarrow 0$  or  $|w_{21}| \rightarrow 0$ , the two solitons decay into a soliton as shown in Fig. 7(d). Those studies are of great help in controlling the oscillation intensity of parabolic solitons so that it can promote theoretical and experimental research on soliton transmission and collision.




**Fig. 3.** Phase shift affects the parallel soliton with oscillations. The parameters are  $\theta_{11} = -1$ ,  $\theta_{12} = 2$ ,  $\theta_{21} = -1$ ,  $\theta_{22} = 2$ ,  $\beta_1(z) = 2e^{-5z^2} + \sin z$  and  $\gamma(z) = 4e^{-0.5z^2}$  with (a):  $w_{11} = 0.47$ ,  $w_{12} = -1.3$ ,  $w_{21} = -0.56$  and  $w_{22} = 1.2$ ; (b):  $w_{11} = -0.84$ ,  $w_{12} = -0.59$ ,  $w_{21} = 0.53$  and  $w_{22} = 0.22$ ; (c):  $w_{11} = 0.5$ ,  $w_{12} = -1.4$ ,  $w_{21} = 0.75$  and  $w_{22} = -0.84$ ; (d):  $w_{11} = -1.3$ ,  $w_{12} = 0.41$ ,  $w_{21} = 0.81$  and  $w_{22} = 0.19$ .




**Fig. 4.** Amplitude comparison between two solitons at  $z = -11$  (red dotted line) and  $z = 11$  (green solid line) in (a): Fig. 3(a); (b): Fig. 3(b); (c): Fig. 3(c); (d): Fig. 3(d).






**Fig. 5.** The parallel soliton amplification in bound states. The parameters are  $\theta_{11} = -1$ ,  $\theta_{12} = 2$ ,  $\theta_{21} = -1$ ,  $\theta_{22} = 2$ ,  $\beta_1(z) = -\frac{0.55}{1+5z^2}$  and  $\gamma(z) = 4e^{-0.5z^2}$  with (a):  $w_{11} = 0.8$ ,  $w_{12} = 0.69$ ,  $w_{21} = -0.78$  and  $w_{22} = -0.41$ ; (b):  $w_{11} = 0.59$ ,  $w_{12} = -0.8$ ,  $w_{21} = -0.38$  and  $w_{22} = -0.55$ ; (c):  $w_{11} = 0.5$ ,  $w_{12} = 0.69$ ,  $w_{21} = -0.94$  and  $w_{22} = 0.59$ ; (d):  $w_{11} = -0.94$ ,  $w_{12} = 0.67$ ,  $w_{21} = 0.63$  and  $w_{22} = -0.63$ .



**Fig. 6.** Amplitude comparison between two solitons at  $z = -16$  (red dotted line) and  $z = 16$  (green solid line) in (a): Fig. 5(a); (b): Fig. 5(b); (c): Fig. 5(c); (d): Fig. 5(d).




**Fig. 7.** The interactions between parabolic solitons with oscillations. The parameters are  $\theta_{11} = -1, \theta_{12} = 2, \theta_{21} = -1, \theta_{22} = 2, \beta_1(z) = -\cos(\arccos z)$  and  $\gamma(z) = \sin 2z$  with (a):  $w_{11} = -0.59, w_{12} = -0.94, w_{21} = 0.83$  and  $w_{22} = -1.9$ ; (b):  $w_{11} = -0.59, w_{12} = 0.016, w_{21} = 0.83$  and  $w_{22} = -1.9$ ; (c):  $w_{11} = -0.59, w_{12} = -0.94, w_{21} = 0.83$  and  $w_{22} = 0.25$ ; (d):  $w_{11} = -0.031, w_{12} = -0.94, w_{21} = 0.83$  and  $w_{22} = -1.9$ .

In order to further explore the effect of  $\beta_1(z)$  and  $\gamma(z)$  on other types of soliton transmissions, one takes  $\beta_1(z) = \arcsin z$  and  $\gamma(z) = \text{arcsinh } z$  as shown in Fig. 8, so that the parabolic solitons can decay into two dromion-like structures. The method of how to adjust the parameters to achieve the suitable attenuation is proposed. The attenuation of solitons is mainly influenced by  $w_{11}, w_{12}$  and  $w_{22}$ . The smaller the  $|w_{11}|$  value, the faster the decay rate becomes. Only when  $w_{12}$  and  $w_{22}$  are positive can the parabolic solitons be attenuated to the dromion-like structures. Further, as  $w_{12}$  and  $w_{22}$  increase, the decay rate becomes faster. Moreover, the distance between two dromion-like structures can be adjusted by  $w_{ij} (i = 1, 2, j = 1, 2)$  as shown in Fig. 8(a) and (b). Their opening

directions are related to the sign of  $A$  in  $\gamma(z) = \text{arcsinh} Az$ . These parametric analyses are of great significance in the development of Bose-Einstein condensation.

**Conclusions**

Two soliton solutions of Eq. (1) were obtained through use of the Hirota method. By analysing the solution, different function forms for  $\beta_1(z)$  and  $\gamma(z)$  were taken to obtain different types of soliton transmission. The effect of the phase shift on the transmission of smooth parallel solitons was discussed when  $\beta_1(z) = 2e^{-0.5z^2}$  and



**Fig. 8.** The parabolic solitons decay into two dromion-like structures. The parameters are  $\theta_{11} = 1, \theta_{12} = 1, \theta_{21} = 1, \theta_{22} = 1, \beta_1(z) = \arcsin z$  and  $\gamma(z) = \text{arcsinh } z$  with (a):  $w_{11} = -0.33, w_{12} = 0.88, w_{21} = 0.14$  and  $w_{22} = 0.36$ ; (b):  $w_{11} = -0.33, w_{12} = 0.88, w_{21} = 0.14$  and  $w_{22} = 0.89$ .

$\gamma(z) = 2\beta_1(z)$ . If one takes  $\beta_1(z) = 2e^{-5z^2} + \sin z$  and  $\gamma(z) = 4e^{-0.5z^2}$ , the influence of the phase shift on parallel solitons was presented. In addition, the soliton amplification in a bound state has been explored when  $\beta_1(z) = -\frac{0.55}{1+5z}$ . When  $\beta_1(z) = -\cos(\arccos z)$  and  $\gamma(z) = \sin 2z$ , parabolic solitons with oscillations are demonstrated. Moreover, parabolic solitons have been reduced to dromion-like structures while  $\beta_1(z) = \arcsin z$  and  $\gamma(z) = \text{arc sinh } z$ . Those presented results are applicable to theoretical analysis and experimental research of optical amplifier, all-optical switches and mode-locked lasers.

### Conflict of interest

*The authors declare they have no conflict of interest.*

### Compliance with Ethics Requirements

*This article does not contain any studies with human or animal subjects.*

### Acknowledgements

This work of Qin Zhou was supported by the National Natural Science Foundation of China (Grant Nos. 11705130 and 1157149). This author was also sponsored by the Chutian Scholar Program of Hubei Government in China. This work of Wenjun Liu was supported by the National Natural Science Foundation of China (Grant No. 11674036), Beijing Youth Top-notch Talent Support Program (Grant No. 2017000026833ZK08), and the Fund of State Key Laboratory of Information Photonics and Optical Communications (Beijing University of Posts and Telecommunications, Grant Nos. IPOC2016ZT04 and IPOC2017ZZ05).

### References

- Wazwaz AM. A new integrable equation combining the modified KdV equation with the negative-order modified KdV equation: multiple soliton solutions and a variety of solitonic solutions. *Wave Random Complex* 2018;28(3):533–43.
- Dong HH, Zhao K, Yang HW, Li YQ. Generalised (2+1)-dimensional super mKdV hierarchy for integrable systems in soliton theory. *E Asian J Appl Math* 2015;5(3):256–72.
- Osman MS, Machado JAT. The dynamical behavior of mixed-type soliton solutions described by (2+1)-dimensional Bogoyavlensky-Konopelchenko equation with variable coefficients. *J Electromagnet Wave* 2018;32(11):1457–64.
- Abdel-Gawad HI, Osman M. On shallow water waves in a medium with time-dependent dispersion and nonlinearity coefficients. *J Adv Res* 2015;6:593–9.
- Qin CY, Tian SF, Wang XB, Zhang TT. On breather waves, rogue waves and solitary waves to a generalized (2+1)-dimensional Camassa-Holm-Kadomtsev-Petviashvili equation. *Commun Nonlinear Sci Numer Simulat* 2018;62:378–85.
- Riaz HWA, Hassan MU. Multi-component semi-discrete coupled dispersionless integrable system, its lax pair and Darboux transformation. *Commun Nonlinear Sci Numer Simulat* 2018;61:71–83.
- Liu ML, Liu WJ, Yan PG, Fang SB, Teng H, Wei ZY. High-power MoTe<sub>2</sub> based passively Q-switched erbium-doped fiber laser. *Chin Opt Lett* 2018;16:020007.
- Cunha AW, Lima ELD, Lima HFD.  $r$ -Almost Newton-Ricci solitons immersed into a Riemannian manifold. *J Math Anal Appl* 2018;464:546–56.
- Yang B, Chen Y. Reductions of Darboux transformations for the PT-symmetric nonlocal Davey-Stewartson equations. *Appl Math Lett* 2018;82:43–9.
- Babalic CN, Carstea AS. Bilinear approach to Kuperschmidt super-KdV type equations. *J Phys* 2018;51:225204.
- Osman MS, Abdel-Gawad HI, El Mahdy MA. Two-layer-atmospheric blocking in a medium with high nonlinearity and lateral dispersion. *Results Phys* 2018;8:1054–60.
- Liu WJ, Liu ML, Lei M, Fang SB, Wei ZY. Titanium selenide saturable absorber mirror for passive Q-switched Er-doped fiber laser. *IEEE J Sel Top Quant* 2018;24:0901005.
- Luo L. The tau-symmetry algebra of the generalized Kaup-Newell hierarchy. *Appl Math Lett* 2018;82:92–7.
- Osman MS. On complex wave solutions governed by the 2D Ginzburg-Landau equation with variable coefficients. *Optik* 2018;156:169–74.
- Sakkaravarthi K, Johnpillai AG, Devi AD, Kanna T, Lakshmanan M. Lie symmetry analysis and group invariant solutions of the nonlinear Helmholtz equation. *Appl Math Comput* 2018;331:457–72.
- Cerdeno DG, Reimtz P, Sakurai K, Tamarit C. B+L violation at colliders and new physics. *J High Energy Phys* 2018;4:076.
- Liu WJ, Liu ML, OuYang YY, Hou HR, Ma GL, Lei M, et al. Tungsten diselenide for mode-locked erbium-doped fiber lasers with short pulse duration. *Nanotechnology* 2018;29:174002.
- Wang YL, Gao YT, Jia SL, Deng GF, Hu WQ. Solitons for a (2+1)-dimensional variable-coefficient Bogoyavlensky-Konopelchenko equation in a fluid. *Mod Phys Lett B* 2017;31:25.
- Kumar A, Dubessy R, Badr T, De Rossi C, de Herve MD, Longchambon L, et al. Producing superfluid circulation states using phase imprinting. *Phys Rev A* 2018;97:4.
- Qadir MI, Zoma T. Symmetric bound states of Josephson vortices in BEC. *Can J Phys* 2018;96:208–12.
- Chen JC, Zhu SD. Residual symmetries and soliton-cnoidal wave interaction solutions for the negative-order Korteweg-de Vries equation. *Appl Math Lett* 2017;73:136–42.
- Baleanu D, Inc M, Yusuf A, Aliyu AI. Optical solitons nonlinear self-adjointness and conservation laws for Kundu-Eckhaus equation. *Chin J Phys* 2018;55:2341–55.
- Yu WT, Yang CY, Liu ML, Zhang YJ, Liu WJ. Interactions of solitons dromion-like structures and butterfly-shaped pulses for variable coefficient nonlinear Schrödinger equation. *Optik* 2018;159:21–30.
- Zhao HQ, Ma WX. Mixed lump-kink solutions to the KP equation. *Comput Math Appl* 2017;74(6):1399–405.
- Apolinar-Fernandez A, Ramos JJ. Numerical solution of the generalized dissipative KdV-RLW-Rosenau equation with a compact method. *Commun Nonlinear Sci* 2018;60:165–83.
- Korkmaz A. Stability satisfied numerical approximates to the non-analytical solutions of the cubic Schrödinger equation. *Appl Math Comput* 2018;331:210–31.
- Feng W, Stanislavova M. On the vortices for the nonlinear Schrödinger equation in higher dimensions. *Philos T R Soc A* 2018;376:20170189.
- Ghanbari B, Inc M. A new generalized exponential rational function method to find exact special solutions for the resonance nonlinear Schrödinger equation. *Eur Phys J Plus* 2018;133:142.
- Osman MS, Korkmaz A, Rezazadeh H, Mirzazadeh M, Eslamif M, Zhou Q. The unified method for conformable time fractional Schrödinger equation with perturbation terms. *Chin J Phys* 2018. doi: <https://doi.org/10.1016/j.cjph.2018.06.009>.
- Cong HZ, Liu JJ, Shi YF, Yuan XP. The stability of full dimensional KAM tori for nonlinear Schrödinger equation. *J Differ Equations* 2018;264:4504–63.
- Correia S. Local Cauchy theory for the nonlinear Schrödinger equation in spaces of infinite mass. *Rev Mat Complut* 2018;31:449–65.
- Kairzhan A, Pelinovsky DE. Nonlinear instability of half-solitons on star graphs. *J Differ Equations* 2018;31:449–65.
- Wang YY, Dai CQ, Xu YQ, Zheng J, Fan Y. Dynamics of nonlocal and localized spatiotemporal solitons for a partially nonlocal nonlinear Schrödinger equation. *Nonlinear Dyn* 2018;92:1261–9.
- Huang ZR, Wang YP, Jia HX, Liu YF. Soliton interaction for a variable-coefficient higher-order nonlinear Schrödinger equation in a dispersion-decreasing fiber. *Opt Laser Technol* 2018;103:151–4.
- Musammil NM, Porsezian K, Nithyanandan K, Subha PA, Dinda PT. Ultrashort dark solitons interactions and nonlinear tunneling in the modified nonlinear Schrödinger equation with variable coefficient. *Opt Laser Technol* 2017;37:11–20.
- Jia XY, Tian B, Liu L, Wu XY, Sun Y. Solitons and breather-to-soliton transitions for an integrable higher-order variable-coefficient nonlinear Schrödinger equation in an optical fiber. *Eur Phys J Plus* 2017;132:488.
- Cai LY, Wang X, Wang L, Li M, Liu Y, Shi YY. Nonautonomous multi-peak solitons and modulation instability for a variable-coefficient nonlinear Schrödinger equation with higher-order effects. *Nonlinear Dyn* 2017;90:2221–30.
- Hasegawa A, Matsumoto M. *Optical solitons in fibers*. Berlin: Springer; 2003.
- Li BQ, Ma YL. Rogue waves for the optical fiber system with variable coefficients. *Optik* 2018;158:177–84.
- Lin XC, Lin WJ, Lei M. Oscillating solitons in nonlinear optics. *Pramana-J Phys* 2016;86:575–80.