



Phase-shift controlling of three solitons in dispersion-decreasing fibers

Suzhi Liu · Qin Zhou · Anjan Biswas ·
Wenjun Liu 

Received: 3 July 2019 / Accepted: 16 August 2019
© Springer Nature B.V. 2019

Abstract Phase-shift controlling can attenuate the interactions between solitons and gives practical advantage in optical communication systems. For the variable-coefficient nonlinear Schrödinger equation, which can be imitated the transmission of solitons in the dispersion-decreasing fiber, analytic three solitons solutions are derived via the Hirota method. Based on the obtained solutions, influences of the second-order dispersion parameters and other related parameters in different function types on the soliton transmission are discussed. Results declare that phase-shift controlling of solitons in dispersion-decreasing fiber can be achieved when the dispersion function is Gaussian one.

S. Liu · W. Liu (✉)
State Key Laboratory of Information Photonics and Optical Communications, and School of Science,
Beijing University of Posts and Telecommunications,
P. O. Box 122, Beijing 100876, China
e-mail: jungliu@bupt.edu.cn

Q. Zhou
School of Electronics and Information Engineering, Wuhan Donghu University, Wuhan 430212,
People's Republic of China

A. Biswas
Department of Physics, Chemistry and Mathematics,
Alabama A&M University, Normal, AL 35762, USA

A. Biswas
Department of Mathematics, King Abdulaziz University,
Jeddah 21589, Saudi Arabia

A. Biswas
Department of Mathematics and Statistics, Tshwane University
of Technology, Pretoria 0008, South Africa

In addition, by adjusting the constraint value, propagation distance of solitons can be further extended. This may be useful in the optical logic devices and ultra-short pulse lasers.

Keywords Soliton · Analytic solution · Dispersion-decreasing fiber · Hirota method

1 Introduction

Optical solitons, which can be stably transmitted by virtue of the offset about the group velocity dispersion (GVD) and self-phase modulation (SPM), have induced substantial interests of researchers in mathematics, physics and optics [1–24]. In particular, the nonlinearity and integrability in the solitons equation are widely concerned [25–28]. Under ideal conditions, solitons can achieve lossless transmission [29]. However, when two- or multi-solitons are transmitted simultaneously in the optical fiber, these solitons can attract each other [29–31]. Due to the existence of solitary interactions, which causes transmission rate of the optical communication system to be seriously degraded, studying how to minimize the interactions becomes one of the meaningful directions [32,33].

Phase-shift controlling, which contributes to an approach for the question mentioned above, has been investigated widely in theory and experiment [34,35]. Ref. [34] has researched large phase shift of solitons in lead glass under strong nonlocal space. Furthermore,

the phase shifts and trajectories during the overtaking interaction of multi-solitons have been analyzed in Ref. [35]. The propagation of three solitons in the dispersion-decreasing fiber (DDF) can be explained by the nonlinear Schrödinger (NLS) equation as follows [36,37]:

$$\phi_\xi - i \frac{D(\xi)}{2} \phi_{\tau\tau} + i\rho(\xi)|\phi|^2\phi = g(\xi)\phi. \tag{1}$$

Here, ϕ is complex function with ξ and τ representing scaled distance and time. $D(\xi)$ is the GVD parameter, $\rho(\xi)$ is the Kerr nonlinear parameter, and $g(\xi)$ is a parameter related to fiber loss or amplification. With regard to Eq. (1), the formation and amplification of solitons have been studied [38]. Interactions between two solitons have been attenuated by phase-shift controlling [36]. Besides, researches have done about how to amplify, reshape, fission and annihilate solitons [39] and investigated the propagation properties of optical solitons in the DDF [40].

Compared with two solitons, the results of three solitons on propagation and interactions are more abundant. In order to better understand the phase-shift characteristics of three solitons in DDF, results of interactions are shown and analyzed under the GVD parameters with different functions in this paper. By setting appropriate dispersion values, phase-shift controlling of solitons in DDF can be better achieved, which is helpful to improve the transmission quality of the signal in optical communication system. In addition, the loss in the medium can be effectively compensated by changing the constraint value produced by $D(\xi)$, $\rho(\xi)$ and $g(\xi)$, so that the propagation distance can be extended.

The structure of the paper is as follows. Section 2 is arranged to get analytic three solitons solutions of Eq. (1) through the Hirota method. Section 3 is reserved for analyzing interactions influences under different value of related parameters and finding suitable value to weaken the interactions among three solitons. Finally, Sect. 4 is used to compose a conclusion for this paper.

2 Bilinear forms and three solitons solutions

Using the Hirota method to introduce the dependent variable transformation:

$$\phi(\xi, \tau) = \frac{h(\xi, \tau)}{f(\xi, \tau)} e^{\int g(\xi) d\xi}, \tag{2}$$

where $h(\xi, \tau)$ is assumed as a complex differentiable function while $f(\xi, \tau)$ is a real one. After calculation, the following bilinear equations of Eq. (1) are obtained:

$$\begin{aligned} (2iD_\xi + D(\xi)D_\tau^2)h \cdot f &= 0, \\ D(\xi)D_\tau^2 f \cdot f + 2\rho(\xi)e^{2\int g(\xi)dx} h \cdot h^* &= 0, \end{aligned} \tag{3}$$

where * denotes complex conjugate. There is a constraint relationship among $D(\xi)$, $\rho(\xi)$ and $g(\xi)$:

$$\rho(\xi) = P \frac{D(\xi)}{e^{2\int g(\xi)d\xi}}. \tag{4}$$

As Hirota bilinear operators, D_ξ and D_τ have the following form:

$$\begin{aligned} D_\xi^m D_\tau^n (a \cdot b) &= \left(\frac{\partial}{\partial \xi} - \frac{\partial}{\partial \xi'} \right)^m \left(\frac{\partial}{\partial \tau} - \frac{\partial}{\partial \tau'} \right)^n \\ &a(\xi, \tau)b(\xi', \tau')|_{\xi'=\xi, \tau'=\tau}, \end{aligned} \tag{5}$$

where both of m and n are non-negative integers, a is the function about ξ and τ , and b is the function about ξ' and τ' . To solve bilinear forms (3), $h(\xi, \tau)$ and $f(\xi, \tau)$ can be expanded as:

$$\begin{aligned} h(\xi, \tau) &= \epsilon h_1(\xi, \tau) + \epsilon^3 h_3(\xi, \tau) + \epsilon^5 h_5(\xi, \tau), \\ f(\xi, \tau) &= 1 + \epsilon^2 f_2(\xi, \tau) + \epsilon^4 f_4(\xi, \tau) + \epsilon^6 f_6(\xi, \tau). \end{aligned} \tag{6}$$

Without affecting the generality, we can define the value of ϵ as 1, and expand the bilinear forms (3) by using expression (6). Three solitons solutions of Eq. (1) are constructed as follows:

$$\phi(\xi, \tau) = \frac{h_1(\xi, \tau) + h_3(\xi, \tau) + h_5(\xi, \tau)}{1 + f_2(\xi, \tau) + f_4(\xi, \tau) + f_6(\xi, \tau)} e^{\int g(\xi) d\xi}, \tag{7}$$

where

$$\begin{aligned} h_1(\xi, \tau) &= e^{\theta_1} + e^{\theta_2} + e^{\theta_3}, \\ f_2(\xi, \tau) &= A_1(\xi)e^{\theta_1^* + \theta_1} + A_2(\xi)e^{\theta_2^* + \theta_1} \\ &\quad + A_3(\xi)e^{\theta_3^* + \theta_1} + A_4(\xi)e^{\theta_1^* + \theta_2} \\ &\quad + A_5(\xi)e^{\theta_2^* + \theta_2} + A_6(\xi)e^{\theta_3^* + \theta_2} \\ &\quad + A_7(\xi)e^{\theta_1^* + \theta_3} + A_8(\xi)e^{\theta_2^* + \theta_3} \\ &\quad + A_9(\xi)e^{\theta_3^* + \theta_3}, \\ h_3(\xi, \tau) &= B_1(\xi)e^{\theta_1^* + \theta_1 + \theta_2} + B_2(\xi)e^{\theta_2^* + \theta_1 + \theta_2} \\ &\quad + B_3(\xi)e^{\theta_3^* + \theta_1 + \theta_2} + B_4(\xi)e^{\theta_1^* + \theta_1 + \theta_3} \end{aligned}$$

$$\begin{aligned}
 & + B_5(\xi)e^{\theta_2^*+\theta_1+\theta_3} + B_6(\xi)e^{\theta_3^*+\theta_1+\theta_3} \\
 & + B_7(\xi)e^{\theta_1^*+\theta_2+\theta_3} + B_8(\xi)e^{\theta_2^*+\theta_2+\theta_3} \\
 & + B_9(\xi)e^{\theta_3^*+\theta_2+\theta_3}, \\
 f_4(\xi, \tau) = & M_1(\xi)e^{\theta_1^*+\theta_2^*+\theta_1+\theta_2} + M_2(\xi)e^{\theta_1^*+\theta_3^*+\theta_1+\theta_2} \\
 & + M_3(\xi)e^{\theta_2^*+\theta_3^*+\theta_1+\theta_2} + M_4(\xi)e^{\theta_1^*+\theta_2^*+\theta_1+\theta_3} \\
 & + M_5(\xi)e^{\theta_1^*+\theta_3^*+\theta_1+\theta_3} + M_6(\xi)e^{\theta_2^*+\theta_3^*+\theta_1+\theta_3} \\
 & + M_7(\xi)e^{\theta_1^*+\theta_2^*+\theta_2+\theta_3} + M_8(\xi)e^{\theta_1^*+\theta_3^*+\theta_2+\theta_3} \\
 & + M_9(\xi)e^{\theta_2^*+\theta_3^*+\theta_2+\theta_3}, \\
 h_5(\xi, \tau) = & N_1(\xi)e^{\theta_1^*+\theta_2^*+\theta_1+\theta_2+\theta_3} + N_2(\xi)e^{\theta_1^*+\theta_3^*+\theta_1+\theta_2+\theta_3} \\
 & + N_3(\xi)e^{\theta_2^*+\theta_3^*+\theta_1+\theta_2+\theta_3}, \\
 f_6(\xi, \tau) = & n(\xi)e^{\theta_1^*+\theta_2^*+\theta_3^*+\theta_1+\theta_2+\theta_3},
 \end{aligned}$$

with

$$\begin{aligned}
 \theta_i & = a_{i1}(\xi) + ia_{i2}(\xi) + (r_{i1} + ir_{i2})\tau + k_{i1} + ik_{i2}, \\
 a_{i1}(\xi) & = - \int r_{i1}r_{i2}D(\xi)d\xi, \\
 a_{i2}(\xi) & = \frac{1}{2} \int (r_{i1}^2 - r_{i2}^2)D(\xi)d\xi (i = 1, 2, 3), \\
 A_1(\xi) & = -\frac{P}{4r_{11}^2}, A_2(\xi) = -\frac{P}{w_{31}^2}, A_3(\xi) = -\frac{P}{w_{32}^2}, \\
 A_4(\xi) & = -\frac{P}{w_{21}^2}, A_5(\xi) = -\frac{P}{4r_{21}^2}, A_6(\xi) = -\frac{P}{w_{33}^2}, \\
 A_7(\xi) & = -\frac{P}{w_{22}^2}, A_8(\xi) = -\frac{P}{w_{23}^2}, A_9(\xi) = -\frac{P}{4r_{31}^2}, \\
 B_1(\xi) & = -\frac{w_{11}^2 P}{4r_{11}^2 w_{21}^2}, B_2(\xi) = -\frac{w_{11}^2 P}{4r_{21}^2 w_{31}^2}, \\
 B_3(\xi) & = -\frac{w_{11}^2 P}{w_{32}^2 w_{33}^2}, \\
 B_4(\xi) & = -\frac{w_{12}^2 P}{4r_{11}^2 w_{22}^2}, B_5(\xi) = -\frac{w_{12}^2 P}{w_{31}^2 w_{23}^2}, \\
 B_6(\xi) & = -\frac{w_{12}^2 P}{4r_{31}^2 w_{32}^2}, \\
 B_7(\xi) & = -\frac{w_{13}^2 P}{w_{21}^2 w_{22}^2}, B_8(\xi) = -\frac{w_{13}^2 P}{4r_{21}^2 w_{23}^2}, \\
 B_9(\xi) & = -\frac{w_{13}^2 P}{4r_{31}^2 w_{33}^2}, \\
 M_1(\xi) & = \frac{J_{11}^2 P^2}{16r_{11}^2 r_{21}^2 J_{12}^2}, M_2(\xi) = \frac{w_{11}^2 w_{42}^2 P^2}{4r_{11}^2 w_{21}^2 w_{32}^2 w_{33}^2}, \\
 M_3(\xi) & = \frac{w_{11}^2 w_{43}^2 P^2}{4r_{21}^2 w_{21}^2 w_{32}^2 w_{33}^2},
 \end{aligned}$$

$$\begin{aligned}
 M_4(\xi) & = \frac{w_{41}^2 w_{32}^2 P^2}{4r_{11}^2 w_{21}^2 w_{22}^2 w_{23}^2}, M_5(\xi) = \frac{J_{21}^2 P^2}{16r_{11}^2 r_{31}^2 J_{22}^2}, \\
 M_6(\xi) & = \frac{w_{12}^2 w_{43}^2 P^2}{4r_{31}^2 w_{31}^2 w_{32}^2 w_{23}^2}, \\
 M_7(\xi) & = \frac{w_{41}^2 w_{13}^2 P^2}{4r_{21}^2 w_{21}^2 w_{22}^2 w_{23}^2}, M_8(\xi) = \frac{w_{13}^2 w_{23}^2 P^2}{4r_{31}^2 w_{21}^2 w_{33}^2 w_{22}^2}, \\
 M_9(\xi) & = \frac{J_{31}^2 P^2}{16r_{21}^2 r_{31}^2 J_{32}^2}, \\
 N_1(\xi) & = \frac{J_{11}^2 w_{12}^2 w_{13}^2 P^2}{16r_{11}^2 r_{21}^2 w_{31}^2 w_{21}^2 w_{22}^2 w_{23}^2}, \\
 N_2(\xi) & = \frac{J_{21}^2 w_{11}^2 w_{13}^2 P^2}{16r_{11}^2 r_{31}^2 w_{21}^2 w_{32}^2 w_{33}^2 w_{22}^2}, \\
 N_3(\xi) & = \frac{J_{31}^2 w_{11}^2 w_{12}^2 P^2}{16r_{21}^2 r_{31}^2 w_{31}^2 w_{32}^2 w_{33}^2 w_{23}^2}, \\
 n(\xi) & = -\frac{J_{11}^2 J_{21}^2 J_{31}^2 P^3}{64r_{11}^2 r_{21}^2 r_{31}^2 w_{31}^2 w_{21}^2 w_{32}^2 w_{33}^2 w_{22}^2 w_{23}^2}, \\
 w_{11} & = r_{11} + ir_{12} - r_{21} - ir_{22}, \\
 w_{12} & = r_{11} + ir_{12} - r_{31} - ir_{32}, \\
 w_{13} & = r_{21} + ir_{22} - r_{31} - ir_{32}, \\
 w_{21} & = r_{11} - ir_{12} + r_{21} + ir_{22}, \\
 w_{22} & = r_{11} - ir_{12} + r_{31} + ir_{32}, \\
 w_{23} & = r_{21} - ir_{22} + r_{31} + ir_{32}, \\
 w_{31} & = r_{11} + ir_{12} + r_{21} - ir_{22}, \\
 w_{32} & = r_{11} + ir_{12} + r_{31} - ir_{32}, \\
 w_{33} & = r_{21} + ir_{22} + r_{31} - ir_{32}, \\
 w_{41} & = r_{11} - ir_{12} - r_{21} + ir_{22}, \\
 w_{42} & = r_{11} - ir_{12} - r_{31} + ir_{32}, \\
 w_{43} & = r_{21} - ir_{22} - r_{31} + ir_{32}, \\
 J_{11} & = (r_{11} - r_{21})^2 + (r_{12} - r_{22})^2, \\
 J_{12} & = (r_{11} + r_{21})^2 + (r_{12} - r_{22})^2, \\
 J_{21} & = (r_{11} - r_{31})^2 + (r_{12} - r_{32})^2, \\
 J_{22} & = (r_{11} + r_{31})^2 + (r_{12} - r_{32})^2, \\
 J_{31} & = (r_{21} - r_{31})^2 + (r_{22} - r_{32})^2, \\
 J_{32} & = (r_{21} + r_{31})^2 + (r_{22} - r_{32})^2.
 \end{aligned}$$




Fig. 1 Interactions among three solitons affected with different profiles of $D(\xi)$. Parameters chosen as: $r_{11} = 2.4, r_{12} = -3.1, r_{21} = -3, r_{22} = 1.8, r_{31} = 3.2, r_{32} = 5, k_{11} = -1, k_{12} = -2, k_{21} = 4, k_{22} = 8, k_{31} = 4, k_{32} = -2, P =$

$-10, g(\xi) = 0.02$ with **a** $D(\xi) = -0.085$; **b** $D(\xi) = -0.07\xi$; **c** $D(\xi) = -0.15\cos(\xi)$; **d** $D(\xi) = -0.07e^{-\xi}$; **e** $D(\xi) = -0.07e^{-\xi^2}$




Fig. 2 Phase shift affected with the different values before ξ^2 . Parameters chosen as: $r_{11} = 2.4, r_{12} = -3.1, r_{21} = -3, r_{22} = 1.8, r_{31} = 3.2, r_{32} = 5, k_{11} = -1, k_{12} = -2, k_{21} =$

$4, k_{22} = 8, k_{31} = 4, k_{32} = -2, P = -10, g(\xi) = 0.02$ with **a** $D(\xi) = -0.057e^{-0.0054\xi^2}$; **b** $D(\xi) = -0.057e^{-0.076\xi^2}$; **c** $D(\xi) = -0.057e^{-0.9\xi^2}$

3 Discussions

For solutions (7) obtained by the above process, we choose different correlative coefficients to observe the interactions among three solitons. Subsequently, the degrees of interactions in different situations are analyzed. Figure 1 indicates interactions among three solitons affected with different profiles of $D(\xi)$ such as the constant one, the linear one, the cosine one, the exponential one and the Gaussian one [41]. In Fig. 1a, we set $D(\xi) = -0.085$. When there is narrow distance between solitons, although influence on the direction of propagation is little, sharp peaks are produced due to the interactions. As shown in Fig. 1b and c, when

$D(\xi) = -0.07\xi$ and $D(\xi) = -0.15\cos(\xi)$, the direction of the soliton propagation is obviously changed and the amplitude of solitons is reduced at the place of interaction especially in Fig. 1b. The attraction between solitons which leads to the appearance of peak can be seen clearly in Fig. 1d. Compared with above profiles, when $D(\xi) = -0.07e^{-\xi^2}$, interactions among three solitons are weakened distinctly because of the phase shift of solitons in Fig. 1e. Results suggest that phase-shift controlling of solitons in DDF can be achieved when the dispersion profile is Gaussian.

In order to present the effect of Gaussian profile on phase shift of solitons, we sequentially change correlative coefficient in Gaussian profile by ten times nearly




Fig. 3 Phase shift affected with the different values before $e^{-0.076\xi^2}$. Parameters chosen as: $r_{11} = 2.4, r_{12} = -3.1, r_{21} = -3, r_{22} = 1.8, r_{31} = 3.2, r_{32} = 5, k_{11} = -1, k_{12} = -2, k_{21} =$

$4, k_{22} = 8, k_{31} = 4, k_{32} = -2, P = -10, g(\xi) = 0.02$ with **a** $D(\xi) = -0.21e^{-0.076\xi^2}$; **b** $D(\xi) = -0.048e^{-0.076\xi^2}$; **c** $D(\xi) = -0.0034e^{-0.076\xi^2}$




Fig. 4 Interactions among three solitons affected with the different values of P . Parameters chosen as: $D(\xi) = -0.05e^{-\xi^2}, r_{11} = 2.4, r_{12} = -3.1, r_{21} = -3, r_{22} = 1.8, r_{31} =$

$3.2, r_{32} = 5, k_{11} = -1, k_{12} = -2, k_{21} = 4, k_{22} = 8, k_{31} = 4, k_{32} = -2, g(\xi) = 0.02$ with **a** $P = -14$; **b** $P = -11$; **c** $P = -8$

as shown in Figs. 2 and 3. With decreasing the value before ξ^2 , the result of the interactions between solitons shows a process from the generation of peaks to the generation of new solitons and then to the independent propagation in Fig. 2. Compared with Fig. 2, Fig. 3 illustrates the same process by increasing the value before $e^{-0.076\xi^2}$. It is found that interactions of solitons can be avoided when the value before ξ^2 is 10 times or larger than before $e^{-\xi^2}$ in Figs. 2 and 3. That means phase-shift controlling can also be realized better by changing the value before ξ^2 or the value before $e^{-\xi^2}$.

Besides, we adjust the value of P in expression (4) as shown in Fig. 4. The attenuation of amplitude becomes alleviative as P increasing, which means the loss in optical fibers can be compensated and the propagation distance of solitons can be improved and extended.

4 Conclusion

Phase-shift controlling of three solitons in DDF has been mainly studied in this paper. Hirota method has been used to obtain the analytic three solitons solutions (7) for Eq. (1). By comparing and analyzing different values of the GVD, the phase-shift controlling has been achieved so that the interactions among three solitons can be weakened effectively by choosing a Gaussian profile. Moreover, we have found that the enhancement of phase-shift controlling can be prompted via decreasing the value before ξ^2 or increasing the value before $e^{-\xi^2}$. In addition, the outcome has been given that the propagation distance of solitons can be prolonged by adjusting the value of P in expression (4). Results maybe can apply in ultra-short pulse lasers and fiber compensation systems.

Acknowledgements The work of Wenjun Liu was supported by the National Natural Science Foundation of China (Grant Nos. 11674036 and 11875008), by the Beijing Youth Top-notch Talent Support Program (Grant No. 2017000026833ZK08) and by the Fund of State Key Laboratory of Information Photonics and Optical Communications (Beijing University of Posts and Telecommunications, Grant No. IPOC2017ZZ05).

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

1. Wazwaz, A.M., El-Tantawy, S.A.: A new integrable (3+1)-dimensional KdV-like model with its multiple-soliton solutions. *Nonlinear Dyn.* **83**(3), 1529–1534 (2016)
2. Wazwaz, A.M.: A study on a two-wave mode Kadomtsev–Petviashvili equation: conditions for multiple soliton solutions to exist. *Math. Methods Appl. Sci.* **40**, 4128–4133 (2017)
3. Wazwaz, A.M.: A two-mode modified KdV equation with multiple soliton solutions. *Appl. Math. Lett.* **70**, 1–6 (2017)
4. Wazwaz, A.M.: Abundant solutions of various physical features for the (2+1)-dimensional modified KdV–Calogero–Bogoyavlenskii–Schiff equation. *Nonlinear Dyn.* **89**, 1727–1732 (2017)
5. Wazwaz, A.M., El-Tantawy, S.A.: Solving the (3+1)-dimensional KP–Boussinesq and BKP–Boussinesq equations by the simplified Hirota’s method. *Nonlinear Dyn.* **88**, 3017–3021 (2017)
6. Wazwaz, A.M.: Multiple soliton solutions and other exact solutions for a two-mode KdV equation. *Math. Methods Appl. Sci.* **40**, 2277–2283 (2017)
7. Nawaz, B., Ali, K., Abbas, S.O., Rizvi, S.T.R., Zhou, Q.: Optical solitons for non-Kerr law nonlinear Schrödinger equation with third and fourth order dispersions. *Chin. J. Phys.* **60**, 133–140 (2019)
8. Rizvi, S.T.R., Ali, K., Hanif, H.: Optical solitons in dual core fibers under various nonlinearities. *Mod. Phys. Lett. B* **33**(17), 1950189 (2019)
9. Rizvi, S.T.R., Afzal, I., Ali, K.: optical solitons for Triki–Biswas equation. *Mod. Phys. Lett. B* **33**, 1950264 (2019)
10. Liu, X.Y., Triki, H., Zhou, Q., Liu, W.J., Biswas, A.: Analytic study on interactions between periodic solitons with controllable parameters. *Nonlinear Dyn.* **94**, 1703–709 (2018)
11. Zhang, Y.J., Yang, C.Y., Yu, W.T., Mirzazadeh, M., Zhou, Q., Liu, W.J.: Interactions of vector anti-dark solitons for the coupled nonlinear Schrödinger equation in inhomogeneous fibers. *Nonlinear Dyn.* **94**, 1351–1360 (2018)
12. Liu, J., Wang, Y.G., Qu, Z.S., Fan, X.W.: 2 μm passive Q-switched mode-locked Tm^{3+} : YAP laser with single-walled carbon nanotube absorber. *Opt. Laser Technol.* **44**(4), 960–962 (2012)
13. Zhang, C., Liu, J., Fan, X.W., Peng, Q.Q., Guo, X.S., Jiang, D.P., Qian, X.B., Su, L.B.: Compact passive Q-switching of a diode-pumped Tm, Y: CaF₂ laser near 2 μm . *Opt. Laser Technol.* **103**, 89–92 (2018)
14. Lin, M.X., Peng, Q.Q., Hou, W., Fan, X.W., Liu, J.: 1.3 μm Q-switched solid-state laser based on few-layer ReS_2 saturable absorber. *Opt. Laser Technol.* **109**, 90–93 (2019)
15. Biswas, A., Yildirim, Y., Yasar, E., Zhou, Q., Moshokoa, S.P., Belic, M.: Optical solitons with differential group delay and four-wave mixing using two integration procedures. *Optik* **167**, 170–188 (2018)
16. Zhang, N., Xia, T.C., Fan, E.G.: A Riemann–Hilbert approach to the Chen–Lee–Liu equation on the half line. *Acta Math. Appl. Sin.* **34**(3), 493–515 (2018)
17. Zhang, N., Xia, T.C., Jin, Q.Y.: N-Fold Darboux transformation of the discrete Ragnisco–Tu system. *Adv. Differ. Equ.* **2018**, 302 (2018)

18. Tao, M.S., Zhang, N., Gao, D.Z., Yang, H.W.: Symmetry analysis for three-dimensional dissipation Rossby waves. *Adv. Differ. Equ.* **2018**, 300 (2018)
19. Gu, J.Y., Zhang, Y., Dong, H.H.: Dynamic behaviors of interaction solutions of (3+1)-dimensional shallow water wave equation. *Comput. Math. Appl.* **76**(6), 1408–1419 (2018)
20. Liu, Y., Dong, H.H., Zhang, Y.: Solutions of a discrete integrable hierarchy by straightening out of its continuous and discrete constrained flows. *Anal. Math. Phys.* **2018**, 1–17 (2018)
21. Guo, M., Zhang, Y., Wang, M., Chen, Y.D., Yang, H.W.: A new ZK-ILW equation for algebraic gravity solitary waves in finite depth stratified atmosphere and the research of squall lines formation mechanism. *Comput. Math. Appl.* **143**, 3589–3603 (2018)
22. Yang, H.W., Chen, X., Guo, M., Chen, Y.D.: A new ZK-BO equation for three-dimensional algebraic Rossby solitary waves and its solution as well as fission property. *Nonlinear Dyn.* **91**, 2019–2032 (2018)
23. Zhao, B.J., Wang, R.Y., Sun, W.J., Yang, H.W.: Combined ZK-mZK equation for Rossby solitary waves with complete Coriolis force and its conservation laws as well as exact solutions. *Adv. Differ. Equ.* **2018**, 42 (2018)
24. Lu, C., Fu, C., Yang, H.W.: Time-fractional generalized Boussinesq equation for Rossby solitary waves with dissipation effect in stratified fluid and conservation laws as well as exact solutions. *Appl. Math. Comput.* **327**, 104–116 (2018)
25. Biswas, A., Rezazadeh, H., Mirzazadeh, M., Eslami, M., Ekici, M., Zhou, Q., Belic, M.: Optical soliton perturbation with Fokas–Lenells equation using three exotic and efficient integration schemes. *Optik* **165**, 288–294 (2018)
26. Triki, H., Biswas, A., Zhou, Q., Mirzazadeh, M., Mahmood, M.F., Moshokoa, S.P., Belic, M.: Solitons in optical metamaterials having parabolic law nonlinearity with detuning effect and Raman scattering. *Optik* **164**, 606–609 (2018)
27. Eslami, M., Mirzazadeh, M.: Optical solitons with Biswas–Milovic equation for power law and dual-power law nonlinearities. *Nonlinear Dyn.* **83**(1–2), 731–738 (2016)
28. Ekici, M., Mirzazadeh, M., Eslami, M.: Solitons and other solutions to Boussinesq equation with power law nonlinearity and dual dispersion. *Nonlinear Dyn.* **84**(2), 669–676 (2016)
29. Li, X.Z., Yan, X., Li, Y.S.: Investigation of multi-soliton, multi-cuspon solutions to the Camassa–Holm equation and their interaction. *Chin. Ann. Math. Ser. B* **33**(2), 225–246 (2012)
30. Sun, K., Tian, B., Liu, W.J., Jiang, Y., Qu, Q.X., Wang, P.: Soliton dynamics and interaction in the Bose–Einstein condensates with harmonic trapping potential and time-varying interatomic interaction. *Nonlinear Dyn.* **67**(1), 165–175 (2012)
31. Askari, A., Javidan, K.: Interaction of noncommutative solitons with defects. *Braz. J. Phys.* **38**(4), 610–614 (2008)
32. Konyukhov, A.I., Dorokhova, M.A., Melnikov, L.A.E., Plastun, A.S.: Controlling the interaction between optical solitons using periodic dispersion variations in an optical fibre. *Quantum Electron.* **45**(11), 1018–1022 (2015)
33. Li, Y.G., She, W.L., Wang, H.C.: Perpendicular all-optical control of interactional optical spatial soliton pair. *Acta Phys. Sin.* **56**(4), 2229–2236 (2007)
34. Shou, Q., Wu, M., Guo, Q.: Large phase shift of (1+1)-dimensional nonlocal spatial solitons in lead glass. *Opt. Commun.* **338**, 133–137 (2015)
35. Roy, K., Ghosh, S.K., Chatterjee, P.: Two-soliton and three-soliton interactions of electron acoustic waves in quantum plasma. *Pramana* **86**(4), 873–883 (2016)
36. Sun, Q.H., Pan, N., Lei, M., Liu, W.J.: Study on phase-shift control in dispersion decreasing fibers. *Acta Phys. Sin.* **63**(15), 150506 (2014)
37. Zheng, H.J., Wu, C., Wang, Z., Yu, H., Liu, S., Li, X.: Propagation characteristics of chirped soliton in periodic distributed amplification systems with variable coefficients. *Optik* **123**(9), 818–822 (2012)
38. Zolotovskii, I.O., Sementsov, D.I.: Formation of the amplification regime of quasi-soliton pulses in waveguides with longitudinally inhomogeneous cross sections. *Opt. Spectrosc.* **102**(4), 594–598 (2007)
39. Yang, C.Y., Li, W.Y., Yu, W.T., Liu, M.L., Zhang, Y.J., Ma, G.L., Lei, M., Liu, W.J.: Amplification, reshaping, fission and annihilation of optical solitons in dispersion-decreasing fiber. *Nonlinear Dyn.* **92**(2), 203–213 (2018)
40. Liu, W.J., Zhang, Y.J., Pang, L.H., Yan, H., Ma, G.L., Lei, M.: Study on the control technology of optical solitons in optical fibers. *Nonlinear Dyn.* **86**(2), 1069–1073 (2016)
41. Pelusi, M.D., Liu, H.F.: Higher order soliton pulse compression in dispersion-decreasing optical fibers. *IEEE J. Quantum Electron.* **33**(8), 1430–1439 (1997)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.